

## An Efficient and One Spot Synthesis of Silica Supported Synthesis of Fluorinated 1,3,4-Thiadiazole Derivative Under Microwave Irradiation

Bharat Dhotre <sup>\*a</sup>, Bhagwan singh Dobhal<sup>b</sup>, Santosh Raut<sup>c</sup>, Amol Kale<sup>a</sup>, Pathan Mohd Arif <sup>c</sup>

<sup>a</sup>Department of Chemistry, Swami Vivekanand Senior College, Mantha, Jalna, Maharashtra, India

<sup>b</sup>Department of Chemistry, Badrinarayan Barwale Mahavidyalaya, Jalna, India

<sup>c</sup> Department of Chemistry, Maulana Azad College, Aurangabad, Maharashtra, India

Corresponding E-mail: [arif7172@rediffmail.com](mailto:arif7172@rediffmail.com)

### Abstract :


An efficient and one pot reaction of fluorinated 1,3,4-Thiadiazole derivatives was synthesis by fluorinated aromatic carboxylic with thiosemicabazide in the presence of phosphorus oxychloride as a catalyst under Microwave irradiation with silica supported. The synthesis compound was characterized by <sup>1</sup>H NMR, IR, and mass spectroscopic techniques. The present approach offers the advantages such as short reaction time, simplicity of the workup procedure, low cost and mild reaction condition.

**Keywords:**1,3,4-Thiadiazole, thiosemicarbazide, Silica supported, Microwave irradiation

### Introduction :

Heterocyclic compounds containing five member rings with nitrogen and sulfur commonly are building blocks, with a possessing large number of interesting biological activities. 1,3,4-thiadiazoles derivatives are five-member with exciting isomeric form of thiadiazoles [1,2]. The important of 1,3,4-thiadiazole is due to its Central Nervous System (CNS) disorders diagnostic ability due the lipo-solubility of sulfur atom. Compounds containing a 1,3,4-thiadiazole template have received significant attention in chemical, medicinal and pharmaceutical research as this structural scaffold is found in a variety of drugs. 1,3,4-thiadiazole and its derivatives are showing immense importance in medicinal chemistry research due to biological activity as well as producing useful intermediates in several organic preparations [3-7]. 1,3,4-Thiadiazoles are heterocyclic scaffolds incorporated in many compounds presenting various pharmacological activities such as antibacterial [8-9], anticonvulsant [10], anti-inflammatory [11-13], antimicrobial [14-17], antiproliferative [18-20], antioxidant [21], antitubercular [22], antitumor [23], anticancer [24-25], antiviral [26]. Similarly, the diagnostic section exhibits a complete investigation of the isomeric form and the anticancer activity of biocaster 1,3,4-thiadiazoles due to therapeutic potential [27-30]. Also fluorinated 1, 3, 4-thiadiazole exhibits more potent antitumor inhibitory activity than doxorubicin [31]. 1,3,4-thiadiazole is central part in several marketed drugs such as acetazolamide, a carbonic anhydrase inhibitor used in treatment of glaucoma or sulfamethizole an antibacterial agent.

### Scheme



## Materials and Methods:







**General:** All the chemical and reagents used were of analytical grade from Aldrich and used without further purification. and the completion of reaction and purity of the synthesized compounds were checked by TLC (0.5 mm thickness) using silica gel-G coated Aluminum plates (Merck). Melting points of the compounds were determined in open capillary tube by digital melting point apparatus and were uncorrected.

### Synthesis of Fluorinated 1,3,4-thiadiazol-2-amine 3 (a-f) under microwave irradiation.

The mixture of fluoro substituted aromatic carboxylic acid (0.01 mole), thiosemicarbazide (0.01 mole), Silica oxide and 5 ml of phosphorus oxychloride were added. The mixture was irradiated under microwave irradiation at 120 °C for 15 min. The completion of the reaction was monitored by TLC. After completion of the reaction the RBF was removed from the oven. The reaction mixture was poured on to crushed ice drop wise with continuous stirring, neutralized by saturated KOH. Then filter, dried and re-crystallized from ethanol.

## Results and Discussion

**Table .1.** Physical data of synthesized compounds 3(a-f) under microwave irradiation

Entry	Compound	Ar	Reaction Time (min)	Yield (%)	M.P.(°C)
1	3a		15	91	<b>222</b>
2	3b		15	90	<b>216</b>
3	3c		15	88	<b>236</b>
4	3d		15	90	<b>155</b>
5	3e		15	90	165
6	3f		15	88	236

An efficient, simple and one pot procedure is reported for the synthesis of fluorinated 1,3,4-Thiadiazole derivative with silica-support in the presence of phosphorus oxychloride microwave radiation technique. 1,3,4-

Thiadiazole derivative are synthesized by fluorinated aromatic carboxylic acid under above of mention condition. The formation of substituted 1,3,4-Thiadiazole derivative were confirmed by recording their IR,  $^1\text{H}$  NMR, and mass spectral data. The compound 5-(2,3,5,6-tetrafluorophenyl)-1,3,4-thiadiazol-2-amine (**3a**) does not shows absorption band at 1700 (carbonyl stretching) broad band at 815 (C-S-C) is present. It shows absorption band at 1095, 1417, 1603 $\text{cm}^{-1}$  is for to C-F, C=C, C=N, stretching respectively.  $^1\text{H}$  NMR of compound **3a** showed broad singlet in the region of  $\delta$ :8.02 which is due to de-shielded caused by the four fluorine proton, the singlet at 8.02 for the  $\text{NH}_2$  of thiadiazole. The mass spectrum of **3a** showed a molecular ion peak at  $m/z$  250 which is in agreement with the molecular formula  $\text{C}_8\text{H}_3\text{F}_4\text{N}_3\text{S}$ . The IR absorption band around 815  $\text{cm}^{-1}$  could be attributed to the -C-S-C functional group. IR spectrum of **3b** showed absorption bands at 3369, 1443, 1596, 1054  $\text{cm}^{-1}$  indicating the presence of  $\text{NH}_2$ , C=C, C=N, C-F groups respectively.  $^1\text{H}$ NMR spectrum peaks due to  $\text{NH}_2$  protons appeared at  $\delta$ :7.46. Peaks for three aromatic proton appeared between 7.20-8.12. Further, LC mass spectrum showed molecular ion peak at  $m/z$  214 which is in conforming the Molecular Structure of **3b**.

## Conclusions

In conclusion, we have reported an efficient and one pot method for the synthesis of fluorinated 1,3,4-Thiadiazole by using fluorinated aromatic acid and thiosemicarbazide in of presence phosphorus oxychloride of under microwave irradiation. This method offers several advantages, including the low of cost, high yields, clean reactions, and short reaction time for the synthesis of one pot synthesis of 1,3,4-thiadiazole. These derivatives have been given a key to more modifications in pharmacophore replacements.

## Data Availability

### 5-(2,4-Difluoro-phenyl)-[1,3,4]thiadiazol-2-ylamine (**3a**)

Yield 85 % m.p.216°C; IR spectrum,  $\nu$ ,  $\text{cm}^{-1}$ : 682(C-S-C stretching), 3369 ( $\text{NH}_2$  stretching.); 1443(C=C Ar stretching), 1596 (C=N stretching), 1054 (C-F stretching);  $^1\text{H}$  NMR spectrum,  $\delta$ , ppm :: 7.20 (s, 1H), 7.41(d, J=8.96, 1H), 7.46(s, 2H), 8.12(d, J=8.96, 1H), (MS:  $m/z$ : 214 (M+H)<sup>+</sup>).

### 5-(3-Chloro-2-fluoro-phenyl)-[1,3,4]thiadiazol-2-ylamine (**3b**)

Yield 84 %, m.p.151°C; IR spectrum $\text{cm}^{-1}$ : 703 (C-S-C stretching), 3385 ( $\text{NH}_2$ stretching.); 1487(C=C Ar stretching), 1607 (C=N stretching), 1009 (C-F stretching);  $^1\text{H}$  NMR spectrum,  $\delta$ , ppm : 7.40 ( dd, J=7.96, 8.4, 1H), 7.52(1H, d, J=7.96, 1H), 8.00(s,2H), 8.16(d, J=8.4, 1H). (MS:  $m/z$ : 230 (M+H)

### 5-(2,3,5,6-Tetrafluoro-phenyl)-[1,3,4]thiadiazol-2-ylamine (**3c**)

Yield 82 %, m.p. 222 °C; IR spectrum,  $\nu$ ,  $\text{cm}^{-1}$ : 793 (C-S-C stretching), 3313 ( $\text{NH}_2$  stretching.); 1413 (C=C Ar stretching), 1595 (C=N stretching), 1030 (C-F stretching);  $^1\text{H}$  NMR spectrum,  $\delta$ , ppm :: 8.02 (s, 1H), 8.01 (s,1H). (MS:  $m/z$ : 250 (M+H)<sup>+</sup>).

## Acknowledgements.

We are thankful to Principal, Maulana Azad College, Aurangabad, and Principal, Swami Vivekanand college Mantha for providing research facilities. We also thank SAIF Punjab University, Chandigarh and SAIF Shillong for providing the spectral and analytical data.

## Conflicts of Interest

Authors do not have any conflict of interest with any person, institution or agency



**Reference:**

1. Shawali, A.S. *Journal of Advanced Research*, 2014, 5, 1–17. <http://dx.doi.org/10.1016/j.jare.2013.01.004>
2. Hamama W. S.; Gouda, M, A; . Abd El-Wahab, M, H.; Zoorob H.H.; *Journal of Heterocyclic Chemistry*, 2014, 51, 1558-1581, DOI 10.1002/jhet.1872
3. Sharma, R.; Sainy, J.; Chaturvedi, S.C. *Acta Pharmaceutica*, 2008, 58, 317–326. DOI 10.2478/v10007-008-0011-6
4. Junyuan T, Jinbing L; Fengyan W, J. Tang et al. *Bio. Chem.* 2016, 69, 29. <http://dx.doi.org/10.1016/j.bioorg.2016.09.007>
5. Pattanayak, P.; Sharma, R.; Sahoo, P. K. *Med. Chem. Res.* 2009, 18, 351. DOI 10.1007/s00044-008-9132-1
6. Rajak, H.; Deshmukh, R.; Aggarwal, N.; Kashaw, S.; Kharya, M. D.; Mishra, P. *Arch. Pharm*, 2009, 342, 453. DOI 10.1007/s00044-008-9132-1
7. S. Maddila a, S. Gorle b, Ch. Sampath c, P. Lavanya J. *Saudi Chem. Soc.* 2016, 20, 306, <http://dx.doi.org/10.1016/j.jscs.2012.11.007>
8. Yousif, E.A.; Majeed, A.S.; Salih, N.A. *Journal of Taibah University for Science*, 2013, 8, 26–30
9. Chandrakantha, B.; Isloor, A.M.; Shetty, P.; Fun, H.K.; Hegde, G. *European Journal of Medicinal Chemistry*, 2014, 71, 316–323.
10. Dogan, H.N.; Duran, A.; Rollas, S. 2002, 10, 2893–2898.
11. Maddila, S.; Gorle, S.; Sampath, C.; Lavanya, P. *Journal of Saudi Chemical Society*, 2016, 20, S306–S312. <http://dx.doi.org/10.1016/j.jscs.2012.11.007>
12. Gadad, A.K.; Palkar, M.B.; Anand, K.; Noolvi, M.N.; Boreddy, T.S.; Wagwade, J. *Bioorganic and Medicinal Chemistry*, 2008, 16, 276–283. doi:10.1016/j.bmc.2007.09.038.
13. Kulkarni, M.V.; Vinay, M.D.; Biradar, S.S.; Rasal, V.P.; Jadhav, V.B. *European Journal of Medicinal Chemistry*, 2008, 43, 1721–1729. <https://doi.org/10.1016/j.ejmech.2007.06.023>
14. Noolvi, M.N.; Patel, H.M.; Kamboj, S. *Arabian Journal of Chemistry*, 2016, 9, 1283–1289. <http://dx.doi.org/10.1016/j.arabjc.2012.02.003>
15. Seelam N.; Shrivastava S.P. *Mccourt, Journal of Saudi Chemical Society* 2016, 20, 33–39 <http://dx.doi.org/10.1016/j.jscs.2012.07.001>
16. Alagawadi, K.R.; Alegaon, S.G. *Arabian Journal of Chemistry*, 2011, 4, 465– 472. doi:10.1016/j.arabjc.2010.07.012
17. Bhardwaj, V.; Noolvi, M.N.; Jalhan, S.; Patel, H.M. *Journal of Saudi, Chemical Society*, 2016, 20, S406–S410. <http://dx.doi.org/10.1016/j.jscs.2012.12.007>
18. Altintop, M.D.; Can, Ö.D.; Özkay, Ü.D.; Kaplancıklı, Z.A. *Molecules*, 2016, 21, 1–10. doi:10.3390/molecules21081004

19. Kumar S.; Gopalakrishnan V. Hegde M.; Rana V.; S. Dhepe S,S; .Bioorganic & Medicinal Chemistry 2014, 24, 4682–4688, <http://dx.doi.org/10.1016/j.bmcl.2014.08.032>
20. Revelant, G.; Gadais, C.; Mathieu, V.; Kirsch, G.; Hesse, S. Bioorganic & Medicinal Chemistry Letters, 2014, 24, 2724–2727. <http://dx.doi.org/10.1016/j.bmcl.2014.04.043>
21. Gür, M; Muğlu H, Çavuş, M,S.; Güder, A.; Sayiner, H, S.; Fatma K. 2014, 43, 36–49. 10.1016/j.molstruc.2016.12.041
22. Ramprasad, J.; Nayak, N.; Dalimba, U.; Yogeewari, P.; Sriram, D.; Peethambar, S.K.; Achur, R.; Kumar, H.S.S. 2015, 95 49-63, <http://dx.doi.org/10.1016/j.ejmech.2015.03.024>
23. Sun, J.; Yang, Y.S.; Li, W.; Zhang, Y. Bin; Wang, X.L.; Tang, J.F.; Zhu, H.L. Bioorganic and Medicinal Chemistry Letters, 2011, 21, 6116–6121. doi:10.1016/j.bmcl.2011.08.039.
24. Kumar, D.; Maruthi Kumar, N.; Chang, K.H.; Shah, K. European Journal of Medicinal Chemistry, 2010, 45, 4664–4668. doi:10.1016/j.ejmech.2010.07.023
25. Polkam, N.; Rayam, P.; Anireddy, J.S.; Yennam, S.; Anantaraju, H.S.; Dharmarajan, S.; Perumal, Y.; Kotapalli, S.S.; Ummanni, R.; Balasubramanian, S. Bioorganic & Medicinal Chemistry Letters, 2015, 25, 1398–1402, <http://dx.doi.org/10.1016/j.bmcl.2015.02.052>
26. Gan, X.; Hu, D.; Chen, Z.; Wang, Y.; Song, B. 2017, 27, 4298–4301. <http://dx.doi.org/10.1016/j.bmcl.2017.08.038>
27. Rzeski, W.; Matysiak, J.; Kandefor-Szerszeń, M. Bioorganic and Medicinal Chemistry, 2007, 15, 3201–3207, doi:10.1016/j.bmc.2007.02.041
28. Matysiak, J.; Nasulewicz, A.; Pelczynska, M.; Switalska, M.; Jaroszewicz, I.; Opolski, A. Eur. J. Med. Chem. 2006, 41, 475.
29. Matysiak, J.; Opolski, A. Bioorg. Med. Chem. 2006, 14, 4483. 10.1016/j.bmc.2006.02.027
30. Zheng, K. B.; He, J.; Zhang, J. Chin. Chem. Lett. 2008, 19, 1281. <https://doi.org/10.1016/j.ccllet.2008.09.021>
31. Xin J. S.; Yu S.; Xing G, D Chin. Chem. Lett , 2011, 22 , 1036. doi:10.1016/j.ccllet.2011.05.012