



On The Gould's Formula for Stirling Numbers of The Second Kind

A. Zúñiga-Segundo ¹, J. López-Bonilla ², S. Vidal-Beltrán ²

¹ ESFM, Instituto Politécnico Nacional, Edif. 9, Dpto. Física, Lindavista 07738, CDMX, México

² ESIME-Zacatenco, Instituto Politécnico Nacional, Edif. 4, 1er. Piso, Col. Lindavista CP 07738, CDMX, México

jlopezb@ipn.mx

Abstract

We present an alternative deduction of the Gould's relation for Stirling numbers of the second kind. Our approach is based in the Nörlund polynomials and in the duality property between the Stirling numbers.

Keywords: Stirling numbers, Duality property, Gould's identity, Nörlund polynomials.

Subject Classification: 11B73, 11B65, 97N70.

Date of Publication: 30-08-2018

Volume: 1 Issue: 2

Journal: MathLAB Journal

Website: <https://purkh.com>



This work is licensed under a Creative Commons Attribution 4.0 International License.



1. Introduction

In Sec. 2 we show the following identity for Stirling numbers of the first kind:

$$S_n^{(n-k)} = \sum_{j=0}^k \binom{n}{k+j} \binom{2k-n}{k-j} S_{k+j}^{(j)}, \quad 1 \leq k \leq n, \quad (1)$$

and in Sec. 3 we exhibit that (1) and the duality property [1-5]:

$$S_{-n}^{(-N)} = (-1)^{N+n} S_N^{[n]}, \quad (2)$$

imply an expression for Stirling numbers of the second kind:

$$S_n^{[n-k]} = \sum_{j=0}^k (-1)^j \binom{n+j-1}{n-k-1} \binom{n+k}{k-j} S_{k+j}^{(j)}, \quad (3)$$

which coincides with the Gould's formula [4, 6]; the relation (3) is the inverse of the Schläfli's identity [7, 8].

2. An identity for Stirling numbers of the first kind

In [9, 10] we have the expression:

$$S_n^{(n-k)} = \binom{n-1}{k} B_k^{(n)}, \quad (4)$$

where $B_n^{(z)}$ are the Nörlund polynomials [11-13] defined via the following generating function:

$$\sum_{n=0}^{\infty} B_n^{(z)} \frac{x^n}{n!} = \left(\frac{x}{e^x - 1} \right)^z; \quad (5)$$

then from (4):

$$S_{k+j}^{(j)} = \binom{k+j-1}{k} B_k^{(k+j)}. \quad (6)$$

On the other hand, in [6, 9] we find the relation:

$$(-1)^k \binom{z}{k} B_k^{(k-z)} = \sum_{j=0}^k \binom{k+j-1}{k} \binom{k-z}{k+j} \binom{k+z}{k-j} B_k^{(k+j)}, \quad (7)$$

that is:

$$(-1)^k \binom{k-n}{k} B_k^{(n)} \stackrel{(6)}{=} \sum_{j=0}^k \binom{n}{k+j} \binom{2k-n}{k-j} S_{k+j}^{(j)}, \quad (8)$$

whose application in (4) implies (1), q.e.d.

3. Gould's formula

In [1] we make the transformation $n \rightarrow -n$ to obtain:

$$S_{-n}^{(-(n+k)} = \sum_{j=0}^k \binom{-n}{k+j} \binom{2k+n}{k-j} S_{k+j}^{(j)}, \quad (9)$$

where we can use (2) and the relation [4, 14]:



$$\binom{-n}{k+j} = (-1)^{k+j} \binom{n+k+j-1}{k+j}, \quad (10)$$

to deduce:

$$S_{n+k}^{[n]} = \sum_{j=1}^k (-1)^j \binom{n+k+j-1}{n-1} \binom{n+2k}{k-j} S_{k+j}^{(j)}, \quad (11)$$

and the change $n \rightarrow n - k$ gives (3), q.e.d.

It is simple to prove that (3) is equivalent to the expression of Gould [4, 6]:

$$S_n^{[n-k]} = (-1)^k \sum_{j=0}^k \binom{k-n}{k+j} \binom{k+n}{k-j} S_{k+j}^{(j)}, \quad (12)$$

because:

$$\binom{k-n}{k+j} = (-1)^{k+j} \binom{n+j-1}{k+j} = (-1)^{k+j} \binom{n+j-1}{n-k-1}. \quad (13)$$

Our procedure shows the usefulness of (2), (10) and (13), that is, of Stirling numbers and binomial coefficients with negative indices [5].

References

1. H. Gupta, Symmetric functions in the theory of integrals numbers, Lucknow University Studies **14**, Allahabad Press (1940)
2. H. Gupta, Selected topics in number theory, Tunbridge Wells, Abacus Press, England (1980)
3. D. E. Knuth, Selected papers on Discrete Mathematics, CSLI Lecture Notes, No. 106 (2003)
4. J. Quaintance, H. W. Gould, Combinatorial identities for Stirling numbers, World Scientific, Singapore (2016)
5. J. López-Bonilla, J. Yaljá Montiel-Pérez, S. Vidal-Beltrán, Stirling numbers with negative indices, World Scientific News **101** (2018) 217-221
6. H. W. Gould, Stirling number representation problems, Proc. Amer. Math. Soc. **11**, No. 3 (1960) 447-451
7. L. Schläfli, Sur les coefficients du développement du produit $1(1+x)(1+2x)\dots(1+(n-1)x$ suivant les puissance ascendantes de x , Crelle's Journal **43** (1852) 1-22
8. J. López-Bonilla, R. López-Vázquez, Harmonic numbers in terms of Stirling numbers of the second kind, Prespacetime Journal **8**, No. 2 (2017) 233-234
9. L. Carlitz, Note on Nörlund polynomials $B_n^{(z)}$, Proc. Amer. Math. Soc. **11**, No. 3 (1960) 452-455
10. I. M. Gessel, On Miki's identity for Bernoulli numbers, J. Number Theory **110**, No. 1 (2005) 75-82
11. N. E. Nörlund, Vorlesungen über differenzenrechnung, Springer-Verlag, Berlin (1924)
12. I. M. Gessel, R. P. Stanley, Stirling polynomials, J. of Combinatorial Theory **A24** (1978) 24-33
13. J. López-Bonilla, S. Yáñez-San Agustín, A. Zaldívar-Sandoval, On the Guo-Qi formula involving Bernoulli and Stirling numbers, Prespacetime Journal **7**, No. 12 (2016) 1672-1673
14. H. M. Srivastava, J. Choi, Zeta and q-zeta functions and associated series and integrals, Elsevier, London (2012)