

## Study of Einstein Solitons and Symmetries of Type $N$ Pure Radiation Fields

M. Danish Siddiqi\* and Shah Alam Siddiqi\*\*

Department of Mathematics, Faculty of Science, Jazan University, Jazan, Kingdom of Saudi Arabia

\*anallintegral@gmail.com , \* msiddiqi@jazanu.edu.sa, \*\*sasiddiqi@jazanu.edu.sa

### Abstract

In the present research paper, we study Einstein solitons with a physical interpretation of the notion of the vector field associated with Einstein solitons. We investigate the geometrical symmetries of Petrov type  $N$  pure radiation fields along the vector field also associated with Einstein solitons.

**Keywords:** Einstein solitons, Petrov type,  $N$  pure radiation, Weyl curvature tensor.

**2010 Mathematics Subject Classification:** 53B50, 53C44, 53C50, 83C02.

### 1. INTRODUCTION

In general theory of relativity, the curvature tensor describing the gravitational field mainly consists of two parts viz, the matter part and the free gravitational part. The interaction between these two parts is described through Bianchi identities. For a given distribution of matter, the construction of gravitational potential satisfying Einstein's field equations is the principal goal of all studies in gravitational physics and this has often been achieved by imposing symmetries on the geometry compatible with the dynamics of the chosen distribution of matter. The geometrical symmetries of the space time are expressible through the vanishing of the Lie derivative of certain tensors with respect to a vector.

The nature of the gravitational radiation from a bounded source is an important physical problem. Even reasonably far from the source, however, twisting type  $N$  solutions of the vacuum field equations are required for an exact description of that radiation. It is well known fact that Petrov type  $N$  solutions of the Einstein vacuum equations are among the most interesting of all empty spacetime metrics [15]. The physical importance, they represent spacetime with gravitational radiation while mathematically they form a class of solution of Einstein equations which should be possible to be determined explicitly.

Recently geometric flows have become important tools in Riemannian geometry and general relativity. In [8] B. List has studied a geometric flow whose fixed points corresponds to static Ricci flat spacetime which is nothing but Ricci flow pullback by a certain diffeomorphism. The association of each Ricci flat spacetime gives notion of local Ricci soliton in one higher dimension. The importance of geometric flow in Riemannian geometry is due to Hamilton who has given the flow equation and B. List generalized Hamilton's equation and extend it to spacetime for static metric [8]. He has given system of flow equations whose fixed points solve the Einstein free-scalar field system [8]. This observation is useful for the correspondence of solutions of system i.e., Ricci soliton and symmetry property of spacetime, that how Riemannian space (or spacetime) with Ricci Soliton deals different kind of symmetry properties.

In the same way like *Ricci solitons* generate self-similar solutions to *Ricci flow*

$$\frac{\partial}{\partial t} g_0 = -2Ric_0 \quad (1.1)$$

In, 2016 G. Catino and L. Mazzeri introduced the notion of Einstein *solitons* [9], which generate self-similar solution to *Einstein flow*

$$\frac{\partial}{\partial t} g_0 = -\left( Ric_0 - \frac{R}{2} g_0 \right) \tag{1.2}$$

The role of symmetries in general theory of relativity has been introduced by Katzin, Levine and Davis in a series of papers ([12], [10]). These symmetries, also known as collineations, were further studied by Ahsan ([1]-[4]), Ahsan and Ali ([5], [6]). The perfect fluid spacetime including electromagnetic field which admit symmetry mapping belonging to the family of contracted Ricci colloneation, have been studied by Norris et al. [13]. The role of geometrical symmetries in the study of fluid spacetime, with an empbasis on conformal collineation has been studied by Duggal [11] and others. The different types of symmetries of Petrov type  $N$  gravitational fields has been subject of interest since last few decades ( cf,[4]).

Motivated by the role of symmetries and Einstein solitons, a study of vector field involved in the definition of Einstein solitons and symmetries of spacetime is made. The main results on the relation between the symmetries of Petrov type  $N$  pure radiation fields and Einstein solitons has been studied.

## 2. Preliminaries

The notion of *Einstein solitons* , which generate self-similar solution to *Einstein flow*.

**(a) Einstein Solitons.** A family  $g_\lambda = \left( \lambda - \frac{R}{2}, x \right)$  of the Riemannian metrics on a  $n$ -dimensional ( $n \geq 3$ ) smooth manifold  $M$  with parameter  $\lambda$  ranging in the time interval  $I \subset \mathbb{R}$  including zero .

Equation (1.2) of Einstein flow for  $g_0 = g(0)$  and the Ricci tensor  $Ric_0$  and curvature tensor  $R$  of the  $g_0$  satisfied. Corresponding to self similar solution of equation (1.2) is the notion of the local Einstein solitons, defined as a metric  $g_0$  satisfying equation

$$-2Ric_0 = L_\xi g_0 + (2\lambda - R)g_0 \tag{2.1}$$

For vector field  $\xi$  on  $V_n$  and a constant  $\lambda$ . The Einstein solitons is said to be steady (static) if  $\lambda = 0$  , shrinking  $\lambda < 0$  and expanding if  $\lambda > 0$ . The metric  $g_0$  is called a gradient Einstein solitons if  $\xi = \nabla \phi$  i.e., gradient of some function  $\phi$ . Schwarzschild metric, Akbar and Woolger [7] have derived the expressions around this notion; while Ali and Ahsan [6] have studied this concept for obtaining the Gaussian curvature of Schwarzschild solitons ad we have.

For  $n$ -dimensional Riemannian manifold equation (2.1) can be written in general as

$$R_{ij} - \frac{1}{2} L_\xi g_{ij} = \left( \lambda - \frac{R}{2} \right) g_{ij} \tag{2.2}$$

So far more than twenty seven different types of collineations have been studied and the literature on such collineations is very large abd still with results of elegance (see [4]). However, here we shall mention only those symmetry assumptions that are required for subsequent investigation and we have

**(b) Motion.** A spacetime is said to admit motion if there exist a vector field  $\xi^t$  such that

$$L_\xi g_{ij} = \xi_{i;j} + \xi_{j;i} = 0 \tag{2.3}$$

equation (2.3) is known as Killing equation and vector  $\xi^t$  is called a Killing vector field [16].

**(c) Conformal Motion .** If

$$L_{\xi}g_{ij} = \sigma g_{ij} \tag{2.4}$$

Where  $\sigma$  is a scalar, then the spacetime is said to admit conformal motion and vector field  $\xi$  is called a conformally Killing vector field.

**(d) Special Conformal Motion.** A spacetime admits if special conformal motion

$$L_{\xi}g_{ij} = \sigma g_{ij}, \quad \sigma_{ij} = 0 \tag{2.5}$$

**(e) Curvature Collineation.** A spacetime admits curvature collineation if there is a vector field  $\xi^i$  such that

$$L_{\xi}R^i_{\ jkl} = 0 \tag{2.6}$$

Where  $R^i_{\ jkl}$  is the Riemannian curvature tensor.

**(f) Ricci Collineation.** A spacetime is said to admit Ricci collineation if there is a vector field  $\xi^i$  such that

$$L_{\xi}R_{ij} = 0 \tag{2.7}$$

Where  $R_{ij}$  is the Ricci tensor.

**(g) Affine Collineation.** If

$$L_{\xi}\Gamma^i_{\ jk} = \xi^i_{\ jk} + R^i_{\ jmk}\xi^m = 0 \tag{2.8}$$

Then spacetime is said to admit an affine collineation.

**(h) Weyl Projective Collineation.** A symmetry property of a spacetime is called Weyl projective collineation if and only if

$$L_{\xi}W^i_{\ jkl} = 0 \quad (n > 2) \tag{2.9}$$

Where  $W^i_{\ jkl}$  is Weyl tensor.

### 3. Main Results

In this section, we shall discuss the role of Einstein solitons in the study of Einstein spaces and Petrov type  $N$  pure radiation fields. In 4-dimension spacetime, the Weyl tensor is related to the Riemannian and Ricci tensors through the equation

$$C_{ijkl} = R_{ijkl} - \frac{1}{2}(g_{ik}R_{jl} + g_{jl}R_{ik} - g_{jk}R_{il} - g_{il}R_{jk}) + \frac{1}{6}(g_{ik}g_{jl} - g_{il}g_{jk})R \tag{3.1}$$

In NP-formalism (cf. [15]), the components of Weyl tensor are expressed by five complex scalars  $\Psi_0, \Psi_1, \Psi_2, \Psi_3$  and  $\Psi_4$ . Through these components the gravitational field has been classified into six categories type *I, II, D, III and O* (cf [15]). The Weyl scalar along with Goldberg-Sachs theorem declares type *N* pure radiation field follow the conditions

$$\Psi_4 = \Psi \neq 0, \quad \Psi_i = 0, \quad i=0, 1, 2, 3 \tag{3.2}$$

and

$$\kappa = \sigma = \varepsilon = 0 \tag{3.3}$$

where  $\kappa, \sigma, \varepsilon$  are the spin-coefficients [15]. Ali and Ahsan [6] have obtained symmetries for Weyl conformal tensor. Using equations (3.1) to (3.3) and the definitions (b)-(c), we can write the following:

**Lemma 3.1.** *In type N pure radiation fields every conformal motion, special conformal motion and homothetic motion, all degenerate to motion.*

From equations (1.4) and (1.5), we have

$$2R_{ij} = L_{\xi} g_{ij} + (2\lambda - R) g_{ij} \tag{3.4}$$

$$= \xi_{i;j} + \xi_{j;i} + (2\lambda - R) g_{ij} \tag{3.5}$$

Contracting this equation with  $g^{ij}$ , we get

$$R = \xi^i{}_{;j} + \left( \lambda - \frac{R}{2} \right) n \tag{3.6}$$

Which can be expressed as

$$div \xi = \nabla_i \xi^i = \left( \frac{(n+2)R}{2} - \lambda n \right) \tag{3.7}$$

where  $R = g^{ij} R_{ij}$  is the scalar curvature. From equations (3.3) and (3.7), we get

$$\left( \frac{(n+1)R}{n} - \frac{R}{2} - R_{ij} \right) = -\frac{1}{2} L_{\xi} g_{ij} + \frac{1}{n} (div \xi) g_{ij} \tag{3.8}$$

Now for  $g_{ij}$  to be Einstein metric i.e.,  $R_{ij} = \sigma g_{ij}$  where  $\sigma$  can be chosen as  $\frac{(n+1)R}{2n}$ , equation (3.3) together with the definition of conformal motion gives the following results:

**Lemma 3.2.** [14] The vector field  $\xi$  associated with Einstein solitons  $(M, g)$  is conformally Killing if and only if  $(M, g)$  is an Einstein manifold of dimension  $(n \geq 3)$ .

Now, using Lemmas 3.1 and 3.2, we can state the following theorem:

**Theorem 3.3.** Type *N* pure radiation field admit motion along a vector field  $\xi$  associated to Einstein solitons  $(M, g)$  if and only if  $M$  is an Einstein space.

For Killing vector field  $\xi$ , equation (1.4) reduces to

$$R_{ij} = \left( \lambda - \frac{R}{2} \right) g_{ij} \tag{3.9}$$

Taking Lie derivative with respect to vector field  $\xi$

$$L_{\xi} R_{ij} = \left( \lambda - \frac{R}{2} \right) L_{\xi} g_{ij} = 0 \tag{3.10}$$

Thus, we have the following theorem:

**Theorem 3.4.** A vector field  $\xi$  associated to Einstein solitons  $(M, g)$  is Ricci collineation vector field in Type  $N$  pure radiation field if  $g$  is Einstein metric.

Taking the Lie derivative of Christoffel symbol  $\Gamma^i_{jk} = \frac{1}{2} g^{il} \left( \frac{\partial g_{jl}}{\partial x^k} + \frac{\partial g_{jk}}{\partial x^l} + \frac{\partial g_{kl}}{\partial x^j} \right)$

along the vector field  $\xi$ , after the calculation we get

$$L_{\xi} \Gamma^i_{jk} = \xi^i_{;jk} + R^i_{jmk} \xi^m \tag{3.11}$$

Now if  $\xi$  is Killing vector field, then

$$\xi^i_{;jk} + R^i_{jmk} \xi^m = 0 \tag{3.12}$$

where 
$$R^h_{ijk} = -\frac{\partial \Gamma^h_{ij}}{\partial x^k} + \frac{\partial \Gamma^h_{ik}}{\partial x^j} - \Gamma^a_{ij} \Gamma^h_{ak} + \Gamma^b_{ik} \Gamma^h_{bj} \tag{3.13}$$

is the Riemannian curvature tensor.

Using (3.11) and (3.12) along with the definition of affine collineation, we can have the following result:

**Theorem 3.5.** Type  $N$  pure radiation field admit affine collineation along a Killing vector field  $\xi$  associated to Einstein solitons  $(M, g)$  if and only if  $M$  is an Einstein space.

By the definition of Lie derivative

$$L_{\xi} R^i_{jkl} = \xi^h R^i_{jkl;h} - R^h_{jkl} \xi^i_{;h} + R^i_{hkl} \xi^h_{;j} + R^i_{jhl} \xi^h_{;k} + R^i_{jkh} \xi^h_{;l} \tag{3.14}$$

Using the definition of Christoffel symbol and Killing vector  $\xi$ , we have

$$L_{\xi} R^i_{jkl} = 0$$

which establishes the curvature collineation, so we have

**Theorem 3.6.** A killing vector field  $\xi$  associated to Einstein solitons  $(M, g)$  is Curvature collineation vector field in type  $N$  pure radiation field if  $g$  is Einstein metric.

The Weyl projective tensor is given by

$$W^i{}_{jkl} = R^i{}_{jkl} - \frac{1}{3}(R_{jk}\delta^i{}_l - R_{jl}\delta^i{}_k)$$

For  $R_{ij} = 0$ ,  $W^i{}_{jkl} = R^i{}_{jkl}$  or  $W_{ijkl} = R_{ijkl}$  (3.15)

From equation (1.8) and (3.15), we can easily write

**Lemma 3.7. [12]** In a Riemannian manifold curvature collineation implies the Weyl projective collineation but converse is true for empty spacetimes.

So, the theorem 3.6 and Lemma 3.7 constitute the following:

**Corollary 3.8.** A Killing vector field  $\xi$  associated to Einstein solitons  $(M, g)$  is Weyl Projective collineation vector field in type  $N$  pure radiation if  $g$  is Einstein metric.

### Conclusions

For Einstein space different kind of symmetry properties for  $N$  pure radiation fields are established with the help of vectr field associated with Einstein solitons. There are other symmetries for type  $N$  which can be obtained through the existence of Killing vectors corresponding to Einstein solitons.

### Acknowledgements

The authors are thankful to the referee for her/his valuable suggestions towards to the improvement of the paper.

### References

1. Ahsan, Z., *Symmetries of the Electromagnetic fields in General Relativity*, Acta Phys. Sincia 4, (1995), 337.
2. Ahsan, Z., *A Symmetry property of the spacetime of general relativity in terms of the space matter tensor*, Brazilin Journal of Phys., 26(3) (1996), 572-576.
3. Ahsan, Z., *Interacting radiation field*, Indian J. Pure App. Maths., 31(2) (2000), 215-225.
4. Ahsan, Z., *On a geometrical symmetry of the spacetime of general relativity*, Bull. Cal. Math. Soc., 97 (3) (2005) 191-200.
5. Ali, M and Ahsan, Z., *Ricci Solitons and Symmetries of spacetime manifold of General relativity*, Glob. J. Adv. Res. Class. Mod. Geom., 1 (2) (2013) 75-84.
6. Ali, M., and Ahsan, Z., *Gravitational field of Schwarzschild soliton*. Arab J. Math. Sci. <http://dx.doi.org/10.1016/j.ajmsc.2013.10.003>.
7. Akbar, M. M., and Woolger, E., *Ricci soliton and Einstein scalar field theory.*, Class. Quantum Grav. 26(2009), 55015.
8. B List, *Evolution of an extended Ricci flow system*, Phd thesis 2005.
9. Catino, G. and Mazzieri, *Gradient Einstein solitons*, Nonlinear Anal. 132 (2016), 66-74.

10. Davis, W. R., Green, L. H. and Norris, L. K., *Relativistic matter fields admitting Ricci collineation and elated conservation laws*, II Nuovo Cimento, 34 B (1976), 256-280.
11. Duggal, K. L., *Relativistic fluids with shear and timelike conformal collineation* J. Math. Phys., 28(1987) 2700-2705.
12. Katzin, G. H. and Levine, J., *Application of Lie derivatives to the symmetries, Geodesic mappings and first integrals in Riemannian spaces*, J. Colloq. Math., 26 (1972) 21-38.
13. Norris, L. K., Green L. H., and Davis, W. R., *Fluid space time including electromagnetic fields admitting symmetry mappings belonging to the family of contracted Ricci collineations*, J. Math. Phys., 18(1977) 1305-1312.
14. Stepanov, S. E. and Shelepova, V. N., *A note on Ricci solitons*, Mathmaticheskije Zametici 86, (3) (2009), 474-477.
15. Stephani, H., Kramer, D., McCallum, M. and Herlt, E., *Exact solutions of Einstein field equations*, Cambridge Univ. Press, Cambridge (2003).
16. Yano, K., *The theory of Lie derivative and its Application*, Vol III, North Holland publishing co. Amsterdam p. Noordhoff L.T.D. Groningen (1957).